

Disease Management Strategies

for Horticultural Crops

B · O · T · A · N · Y



PLANT · PATHOLOGY

www.btny.purdue.edu



Fungicide Rotations for Nursery, Greenhouse, and Landscape Professionals

Janna Beckerman

Department of Botany and Plant Pathology, Purdue University

Fungicides are important tools for managing ornamental plant diseases. There are many different fungicides and numerous methods of classifying them. This publication examines how fungicides are classified and recommends management practices to prevent fungi from developing resistance to these products.

Fungicide Class

One way to classify fungicides is by their chemical structures or **modes of action** — the specific ways the fungicides affect a fungus. Fungicides that share a common mode of action belong to the same **fungicide class** (sometimes referred to as a **fungicide family**). Unfortunately, if a fungus is resistant to a specific fungicide, it is usually resistant to all the fungicides within that fungicide class.

Target Site

Fungicides are also characterized by their specificity.

Site-specific fungicides react with one very specific, very important biochemical process, called the **target site**. For example, a fungicide target site could be the specific proteins involved in cell wall biosynthesis, RNA biosynthesis, or cell division. Site-specific fungicides target these specific processes, which prevents the fungus from growing and ultimately causes its death.

Multi-site fungicides have multiple modes of action, so they affect multiple target sites, and simultaneously interfere with numerous metabolic processes of the fungus.

Fungicide resistance occurs when a fungus develops a genetic mutation at the target site that reduces its sensitivity to a specific fungicide. Because they affect multiple target sites, multi-site fungicides have a very low risk of causing fungicide resistance because it is highly unlikely for a fungus to simultaneously develop all of the mutations necessary for resistance.

Site-specific fungicides, however, have a much higher risk of causing resistance because a single genetic mutation at the target site can change a fungus' biochemical process so that it can still perform the needed biological function (cell division, membrane biosynthesis, respiration). The result is a fungus strain that is less susceptible or no longer susceptible to the site-specific fungicide.

If a single fungicide continues to be used, the fungicide-sensitive portion of the population is suppressed over time, and only the fungicide-resistant portion of the

Purdue Extension

Knowledge to Go

1-888-EXT-INFO

population remains, which goes on to reproduce and make up the majority of the population. Eventually, the fungicide is ineffective because this majority of the fungal population is no longer susceptible to it.

Minimizing Resistance

To minimize the possibility of fungicide resistance from occurring, implement a comprehensive management strategy before resistance develops. Some key tactics to include in your management strategy include:

1. Follow good plant health practices.

Using disease-resistant cultivars, following proper planting and fertilization techniques, and sanitizing equipment reduce the reliance on fungicides, thereby reducing the risk of their over-use and the development of resistant populations.

2. Use the recommended doses as stated on fungicide labels.

Many fungicides have been extensively tested to identify the optimal rate. Cutting the rate results in a sublethal dose that is not only ineffective for disease management, but increases the risk of resistance.

3. Minimize the number of fungicide treatments per season, and apply only when necessary.

Excessive use of site-specific fungicides increases the likelihood of resistance. By reducing the number of site-specific fungicide applications, you reduce the likelihood of resistance development.

4. Do not rely solely on one fungicide with a site-specific mode of action.

Use a diversity of fungicides with different modes of action that provide broad-spectrum disease control. There is no single, best fungicide. There are, however, multiple fungicides with different efficacies for different diseases. Many single-site fungicides are highly effective by themselves, but you should tank-mix them with another fungicide from a different family, or rotate or alternate multiple fungicides to reduce the risk of resistance. The important thing to remember is that you should avoid consecutive applications of site-specific fungicides.

Tank-mixing and Rotating

There are two tactics that can reduce the risk of fungicide-resistant disease populations: tank-mixing and rotating fungicides.

As the name suggests, **tank-mixing** consists of mixing a fungicide with a high resistance risk with another fungicide with a low or negligible resistance risk (Table 1).

Rotating fungicides involves alternating products that have different modes of action so that you avoid back-to-back treatments with any one site-specific fungicide.

Tank-mixing and rotating are important for two reasons. First, both practices limit the amount of time fungi are exposed to any one product. Second, other fungicides could potentially suppress any resistant populations before they have a chance to reproduce.

Selecting the proper tank mix or rotation partners in a fungicide resistance management program is critical. To develop an effective tank mix or rotation:

- Use fungicides with different Group Codes (Table 1), which denote different fungicide families. Remember, fungicides with different trade names can belong to the same chemical family!
- Always partner site-specific products with a multi-site inhibitor fungicide (Group Code M).
- Carefully read fungicide labels to determine if any fungicides cannot be mixed or rotated together.

Fungicide rotations can be simple or complex, depending on the problem and the pathogen that is causing it. For example, you may control a Septoria, Myrothecium, or Alternaria leaf spot by rotating a fludioxonil-based fungicide (Group Code 12) with a chlorothalonil-based fungicide (Group code M), thereby minimizing the risk of resistance in the Group Code 12 fungicide.

Other diseases may require more elaborate rotations. For example, downy mildew on snapdragon or lamium may require:

- A dimethomorph-based fungicide (Group Code 40).
- And then, mancozeb or copper (both Group M).
- And then, a phosphorous acid-based fungicide (Group Code M(33)).
- And then, mancozeb or copper again (Group M).

Furthermore, this rotation could only be repeated twice with the Group 40 fungicide — it is limited to two applications unless it is tank-mixed.

The bottom line is that tank-mixing or rotating fungicides reduces the possibility of resistance development. This is important as the labels on most newer, site-specific fungicides have strict use recommendations to minimize the risk of fungicide resistance, and protect the long-term efficacy of the product. By carefully following these recommendations, and using fungicides with different group codes, diseases and fungicide resistance can be carefully and effectively managed.

Table 1. Fungicides Labeled for Use on Ornamentals. This table provides the common and trade names of selected fungicides currently registered in the United States for use on ornamentals. It also provides the group code, major fungicide families and chemistries within these groups, and the risk of resistance developing due to using these fungicides. Products set in italics indicate that they are prohibited from greenhouse use. Products set in bold indicate that they are for greenhouse use only.

Group Code ¹	Fungicide Family ² or Class	Common Name	Example Trade Name	Risk of Resistance ³
1	benzimidazole or MBC	thiophanate-methyl	3336, [®] Cleary's 3336 [®]	high
2	dicarboximide	iprodione	Chipco 26GT, [®] Iprodione Pro 2SE [®]	medium to high
3	demethylation inhibitor (DMI)	triflumizole	Procure, [®] Terraguard [®]	medium
		triforine	Funginex, [®] Saprol [®]	
		myclobutanil	Eagle, [®] Systhane [®]	
		tebuconazole	Folicur [®]	
		<i>Banner Maxx, Propiconazole</i>	<i>Banner Maxx,[®] Propiconazole[®]</i>	
		bayleton	Bayleton [®]	
4	phenylamide (PA)	mefenoxam	Subdue Maxx [®]	high
7	carboximides	flutalonil	Contrast [®]	medium
		boscalid	Mix partner of Pristine [®]	
11	quinone outside inhibitor (QoI) — strobilurins	kresoxim-methyl	Cygnus[®]	high
		azoxystrobin	Heritage [®]	
		trifloxystrobin	Compass [®]	
		pyraclostrobin	Insignia [®]	
12	phenylpyrrole (PP)	fludioxonil	Medallion [®]	low to medium
14	aromatic hydrocarbons	dicloran	Botran 70 [®]	low
		etridiazole	Truban, [®] Terrazole [®]	
17	hydroxyanilide	fenhexamid	Decree[®]	low to medium
18	antibiotic streptomycetes	streptomycin	Agri-Mycin, [®] Agri-Step [®]	high
19	polyoxin	polyoxin D	Endorse [®]	medium
21(P) ⁴	host plant defense inducers, systemic acquired resistance (SAR)	acibenzolar-S- methyl	Actigard [®]	low
		harpin	Messenger [®]	
M (33)	multi-site activity phosphonate	fosetyl-aluminum	Aliette [®]	low
		phosphorous acid	Alude, [®] BioPhos [®]	
40	cinnamic acid	dimethomorph	Stature DM, [®] Stature SC [®]	low to medium
M	multi-site activity inorganics	copper	Kocide, [®] Champ [®]	low
M	multi-site activity inorganics	sulfur	Microthiol Disperss, [®] Sulfur	low
M	multi-site activity dithiocarbamate	mancozeb, maneb, dimethyldithio-carbamate	Mancozeb, [®] Maneb, [®] Thiram [®]	low
M	multi-site activity chloroalkythios	captan	Captan [®]	low
M	multi-site activity chloronitrile	chlorothalonil	Bravo, [®] Daconil 2787 [®]	low
U(28)	unknown carbamate	propamocarb	Banol [®]	low to medium

¹ FRAC code is listed in parentheses under the EPA Group code when the codes differ. Neither system includes biofungicides.

² For the sake of consistency, group codes, fungicide classes, fungicide names, and abbreviations are those used by the Fungicide Resistance Action Committee (FRAC) and by the EPA Office of Pesticide Programs. This program is part of the pesticide classification system developed to assist growers in resistance management. Only fungicides registered in the USA are included.

³ Resistance risk is considered high when resistance has already been reported, or a specific, single gene mode of resistance is known within a few years under commercial use; Medium risk is associated with less frequent resistance development, or when more than one gene is involved, and the risk is considered low when the fungicide has multi-site activity. Entries in this column were assigned by FRAC (www.frac.info). Labels for fungicides registered in the USA are accessible at www.greenbook.net and www.cdms.net/manuf/manuf.asp.

⁴ Although similarly described, the modes of action are different.

References

McGrath, M.T. 2004. What are Fungicides? *The Plant Health Instructor*. DOI: 10.1094/PHI-I-2004-0825-01.

Thomson, W.T. 2000. Agricultural Chemicals: Book IV Fungicides. Thomson Publications.

Fungicide Resistance Action Committee (FRAC) code list.
www.frac.info/frac/publication/anhang/FRAC_Code_List_2007_web.pdf

Reference to products in this publication is not intended to be an endorsement to the exclusion of others that may be similar. Persons using such products assume responsibility for their use in accordance with current directions of the manufacturer.

For other publications in the Disease Management Strategies series, visit the Purdue Extension Education Store, www.extension.purdue.edu/new.

